



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

NOTE BY THE EDITOR.—As it has been objected that, in the equation we gave, on page 143, (No. 5) of the parallel curve to an ellipse, our result was not satisfactory because the equation involved explicitly only one of the coordinates; we therefore add, as a supplement to that note, the following:

By similar triangles (see fig. on p. 143) we have

$$y - y' : x - x' :: y : ON'. \quad . \quad . \quad . \quad . \quad (1')$$

Also, from the differential triangle,

$$(dx^2 + dy^2)^{\frac{1}{2}} : dy :: N : ON' \quad . \quad . \quad . \quad . \quad (2')$$

From (1') and (2') we get

$$N = \frac{(x - x')y}{y - y'} \sqrt{1 + \frac{dx^2}{dy^2}} = \frac{(x - x')y}{y - y'} \sqrt{1 + \frac{a^4 - a^2x^2}{b^2x^2}}.$$

Writing  $q$  for  $N' \div (N' + c)$ , we get by reduction,

$$x' = x - \frac{N(1 - q)bx}{\sqrt{[a^4 - (a^2 - b^2)x^2]}}.$$

Substituting for  $y$ , in the equation of the ellipse, its value from (2), (p. 143)

we have  $x = \frac{a}{b} \sqrt{(b^2 - q^2y'^2)}$ ; and putting this value of  $x$  for  $x$ , in the

last equation above, and for  $N'$ , as involved in  $q$ , and  $N$ , their values in functions of  $y'$  from the equation given on page 143, we obtain an equation involving only as variables the coordinates  $x'$  and  $y'$ .

---

### PROBLEMS.

---

92. BY A. W. MASON, CEDAR FALLS, IOWA.—A balloon is ascending vertically with a given velocity  $v$ , and a body is let fall from it, which touches the ground in  $t$  seconds; find the height of the balloon at the moment the body is let fall from it.

93. BY PROF. J. SCHEFFER.—To construct a triangle if the three radii of the circles, which touch the three sides externally, be given.

94. BY PROF. W. W. JOHNSON.—One side of a quadrilateral, whose four sides are given in length, is fixed: find the equation of the locus of the middle point of the opposite side, in rectangular coordinates.

95. BY DR. A. B. NELSON, DANVILLE, KY.—Prove, otherwise than by the Integral Calculus that

$$\frac{\pi}{2} - \sin^{-1}e = 2 \tan^{-1} \left( \frac{1 - e}{1 + e} \right)^{\frac{1}{2}}.$$